

# MATH 208, Fall 2011

## Manifolds I

### Final

1. Consider two vector fields  $v(x) = v_0 + A(x)$  and  $w(x) = w_0 + B(x)$  on  $\mathbb{R}^n$ , where  $v_0$  and  $w_0$  are constant vectors, and  $A$  and  $B$  are linear maps  $\mathbb{R}^n \rightarrow \mathbb{R}^n$ .

- a. Find the flow  $\varphi^t$  of  $v$ .
- b. Find the bracket  $[v, w]$ .

2. Let  $\Sigma = \Sigma_g \setminus D^2$ , where  $\Sigma_g$  is the sphere with  $g$  handles and  $D^2$  is a disk. Show that  $\Sigma$  admits an immersion into  $\mathbb{R}^2$ . (It is sufficient to sketch a figure or a series of figures as a solution.)

3. Consider the map  $\tilde{F}: S^2 \rightarrow \mathbb{R}^4$  given by

$$\tilde{F}(x, y, z) = (x^2 - y^2, xy, xz, yz).$$

Prove that  $\tilde{F}$  gives rise to a smooth embedding  $F: \mathbb{R}P^2 \rightarrow \mathbb{R}^4$ .

4. Let  $M$  be a submanifold in  $N \times P$ , where  $N$  and  $P$  are arbitrary manifolds. Show that for almost all  $y \in P$  the intersection  $M \cap (N \times \{y\})$  is a smooth submanifold in  $N \times \{y\} = N$ .

5. Two smooth submanifolds  $M_0$  and  $M_1$  of  $N$  are said to be transverse if  $T_p N = T_p M_0 + T_p M_1$  for all  $p \in M_0 \cap M_1$ . Prove that  $M_0 \cap M_1$  is a smooth submanifold of  $N$  of dimension  $\dim M_0 + \dim M_1 - \dim N$ , whenever  $M_0$  and  $M_1$  are transverse.

6. Denote by  $M_n$  the vector space of real  $n \times n$  matrices and let  $P_n$  be the vector space of real symmetric  $n \times n$  matrices. Consider the map  $F: M_n \rightarrow P_n$  given by  $F(A) = AA^T$ .

- a. Show that  $DF_I(X) = X + X^T$  and that  $I \in M_n$  is a regular point of  $F$ . (Would  $I$  still be a regular point if we replaced the target space  $P_n$  by  $M_n$ ?)
- b. Show that  $I \in P_n$  is a regular value of  $F$ . Thus the orthogonal group  $O(n) = \{A \in M_n \mid AA^T = I\}$  is a smooth submanifold of  $M_n$ . Find  $\dim O(n)$ .
- c. Show that  $T_I O(n)$  is the space of all skew-symmetric matrices, i.e.,  $T_I O(n) = \{X \in M_n \mid X + X^T = 0\}$ . (This space is usually denoted by  $so(n)$ .)

7. Let  $v$  be a vector field on  $M$ . Denote by  $\varphi^t$  the (local) flow of  $v$ .
- Let  $F: M \rightarrow N$  be a diffeomorphism. Show that  $F\varphi^tF^{-1}$  is the (local) flow of  $F_*v$ . (Hint: prove that  $F(\varphi^t(F^{-1}(q)))$  is the integral curve of  $F_*v$  through  $q \in N$ .)
  - Let  $F: M \rightarrow M$  be a diffeomorphism. Prove that  $F_*v = v$  if and only if  $F\varphi^t = \varphi^tF$  for all  $t$ .
8. Let  $M$  be a connected manifold without boundary. Show that for any two points  $p$  and  $q$  in  $M$ , there exists a diffeomorphism  $F: M \rightarrow M$  such that  $F(p) = q$ . In other words, the group of diffeomorphisms of  $M$  acts transitively on  $M$ . (Hint: First reduce the problem to the case where  $p$  and  $q$  are contained in a relatively compact coordinate neighborhood  $U$  diffeomorphic to an open ball. Then show that there exists a vector field  $v$  on  $M$ , vanishing outside  $U$  and hence complete, such that  $\varphi^1(p) = q$ , where  $\varphi^t$  is the flow of  $v$ .)