

MATH 209, MANIFOLDS II, WINTER 2009

Homework Assignment I: One-forms and integration,
due Tuesday 1/21

1. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $y = x^3$. Calculate $f_*(\partial/\partial x)$ and $f^* dy$. Is $f_*(\partial/\partial x)$ smooth?

2. Let f be a smooth function on a manifold M . Show that $df = f^* dy$, where y is the natural coordinate on \mathbb{R} . Furthermore, let (x_1, \dots, x_n) be a system of local coordinates on M . Prove that in local coordinates

$$df = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i.$$

Remark. Here one has to be careful with definitions. In this problem we define the one-form df by $df(v) = L_v f$. The one-forms dx_1, \dots, dx_n are defined as the basis in $T_x^* M$ dual to the basis $\partial/\partial x_1, \dots, \partial/\partial x_n$ in $T_x M$. (Note that alternatively one could define dx_i by $dx_i(v) = L_v x_i$, where x_i is now thought of as a function defined on an open subset of M . It follows from the assertion of the problem that this definition is equivalent to the one through the dual basis.)

3. Let $\alpha = x dy$ on the plane \mathbb{R}^2 with coordinates (x, y) . Denote by S_R^1 the circle of radius $R > 0$ centered at the origin, oriented counter clockwise. Evaluate

$$\int_{S_R^1} \alpha.$$

Conclude from the result that α is not exact, i.e., there exists no function f such that $\alpha = df$. More generally, let γ be a closed simple curve in \mathbb{R}^2 . What is the geometrical meaning of $\int_{\gamma} x dy$?

4. Let α be a one-form on a smooth connected manifold M . Recall that α is said to be exact if there exists a function f such that $\alpha = df$. Prove that

- α is exact $\iff \int_{\gamma} \alpha = 0$ for any closed curve γ .

Remark. The part \implies), which we did in class, is here only for the sake of completeness. You do not need to redo it. Here is a hint for part \impliedby). The goal is to construct f such that $\alpha = df$. Fix a reference point $x_0 \in M$. For $x \in M$, set $f(x) = \int_{\eta} \alpha$, where η is curve connecting x_0 to x . First show that the function f is well defined, i.e., $f(x)$ is really independent of the curve η connecting x_0 and x , using the assumption that $\int_{\gamma} \alpha = 0$. Then prove that $df = \alpha$. (For instance, one can use local coordinates to this end.)

5. Problem 6-7 (Chapter 6) on page 152 of the textbook.