

MATH 209, MANIFOLDS II, WINTER 2011

Final

Throughout the exam all manifolds, maps, and homotopies are assumed to be smooth.

1. Let  $D$  be a closed two-dimensional disk.

- (a) Show that there is no smooth map  $f: D \rightarrow \partial D$  such that  $f|_{\partial D} = id$ .
- (b) Let  $f: D \rightarrow \mathbb{R}^2$  be a smooth map such that the origin  $p = (0, 0)$  is not in  $f(\partial D)$  and the winding number  $W(p, f|_{\partial D}) \neq 0$ . Prove that  $f^{-1}(p) \neq \emptyset$ .

2. Show that  $S^{2n}$  with  $n > 1$  does not admit a symplectic structure.

3. Problem 15-7 on page 408 of the textbook. Prove, as a consequence, that  $H^1(\Sigma_g) = \mathbb{R}^{2g}$ , where  $\Sigma_g$  is the sphere with  $g$  handles.

4. Let  $U$  and  $V$  be open subsets of a manifold  $M$  and let  $\alpha$  be a differential form on  $U \cap V$ . Show that there exist differential forms  $\beta$  on  $U$  and  $\gamma$  on  $V$  such that  $\alpha = \beta|_{U \cap V} - \gamma|_{U \cap V}$ . (This problem completes the construction of the Mayer-Vietoris exact sequence.)

5. The objective of this problem is to give a direct proof of Poincaré's lemma. Let  $v$  be a vector field on a manifold  $M$ . Assume that the flow  $\varphi_t$  of  $v$  is defined for all  $t \in \mathbb{R}$ .

- (a) Prove that  $i_v \varphi_t^* \alpha = \varphi_t^* i_v \alpha$  for any differential form  $\alpha$ .
- (b) Prove that

$$\frac{d}{dt} \varphi_t^* \alpha = L_v \varphi_t^* \alpha = \varphi_t^* L_v \alpha$$

for all  $t \in \mathbb{R}$  and any differential form  $\alpha$ .

Let now  $M = \mathbb{R}^n$  and  $v$  be given by  $v(x) = -x$ ,  $x \in \mathbb{R}^n$ . (Here we identify  $T_x \mathbb{R}^n$  and  $\mathbb{R}^n$ .) Let  $k \geq 1$ . Consider the linear operator  $\Phi_k: \Omega^k(\mathbb{R}^n) \rightarrow \Omega^{k-1}(\mathbb{R}^n)$  defined by

$$(\Phi_k \alpha)(w_1, \dots, w_{k-1}) = - \int_0^\infty (\varphi_t^* i_v \alpha)(w_1, \dots, w_{k-1}) dt$$

for a  $k$ -form  $\alpha$  on  $\mathbb{R}^n$ .

- (c) Prove that the improper integral in the definition of  $\Phi_k \alpha$  converges.
- (d) Complete the proof of Poincaré's lemma by showing that  $d\Phi_k \alpha + \Phi_{k+1} d\alpha = \alpha$  for any  $k$ -form  $\alpha$  on  $\mathbb{R}^n$ . (Thus, if  $\alpha$  is closed,  $\alpha = d\Phi_k \alpha$  and, hence,  $\alpha$  is exact.) Hint: Use Part (b) and Cartan's formula!

6\*. (Hopf invariant) The goal of this problem is to introduce a homotopy invariant of maps  $f: S^3 \rightarrow S^2$ , the so-called Hopf invariant, and to establish some of its properties.

- (a) Let  $\omega \in \Omega^2(S^2)$  be an arbitrary form with  $\int_{S^2} \omega = 1$ . Since  $H^2(S^3) = 0$ , the pull back  $f^* \omega$  is exact. Define the Hopf invariant of  $f$  by  $H(f) = \int_{S^3} \lambda \wedge f^* \omega$ , where  $\lambda$  is a primitive of  $f^* \omega$ , i.e.,  $d\lambda = f^* \omega$ . Prove that  $H(f)$  is well-defined, i.e., independent of the choice of  $\lambda$  when  $\omega$  is fixed and, furthermore, independent of the choice of  $\omega$ .
- (b) Evaluate the Hopf invariant for the Hopf fibration  $S^3 \rightarrow S^2$ .
- (c) Prove that  $H(f)$  is homotopy invariant.

Remark. One can also show that  $H(f) \in \mathbb{Z}$  for all  $f$  and  $f_0 \sim f_1$  iff  $H(f_0) = H(f_1)$ .