

MATH 209, MANIFOLDS II, WINTER 2023

Homework Assignment V: Orientations and integration

1. Let  $F: \mathbb{C} \rightarrow \mathbb{C}$  be a holomorphic function. Show that  $F$  is necessarily orientation preserving at its regular points, i.e.,  $F^*dx \wedge dy = f dx \wedge dy$  with  $f \geq 0$ .

2. Let  $N$  be a hypersurface in  $M$  and let  $\omega$  be a volume form on  $M$ , i.e., a non-vanishing top-degree form.

(a) Let  $v$  be a vector field nowhere tangent to  $N$ . Prove that  $i_v\omega|_N$  is a volume form on  $N$ .

(b) Prove that  $i_v\omega|_N = i_w\omega|_N$  if  $v - w$  is tangent to  $N$ .

Remark. Assume that the hypersurface  $N$  is the boundary of  $M$ . Then the construction of Part (a) gives an alternative description of the orientation induced on  $N$ . Indeed, let  $v$  point outward and let an orientation of  $M$  be determined by  $\omega$ . Then the induced orientation of  $N = \partial M$  is determined by  $i_v\omega$  and is well defined. Note also that in both (a) and (b) it suffices to have  $v$  defined only along  $N$ .

3. Let  $\omega \in \Omega^2(M)$  and  $u: D \rightarrow M$  be a smooth map, where  $D$  is a compact domain in  $\mathbb{R}^2$  with smooth boundary. Prove that in the notation introduced in class

$$u^*\omega = \omega \left( \frac{\partial u}{\partial t}, \frac{\partial u}{\partial s} \right) dt \wedge ds$$

where  $t$  and  $s$  are the coordinates on the domain  $D$  of  $u$  and, as a consequence,

$$\int_u \omega = \iint_D \omega \left( \frac{\partial u}{\partial t}, \frac{\partial u}{\partial s} \right) dt ds.$$

4. In the setting of the previous problem with  $M = \mathbb{R}^3$ , let us think of  $u = (u_1, u_2, u_3): D \rightarrow \mathbb{R}^3$  as a parametrization of a surface by  $D \subset \mathbb{R}^2$  in the sense of vector calculus. We assume for the sake of simplicity that  $u = u(D)$  is an embedded surface with boundary with the orientation inherited from that of  $D$ . Then

$$\frac{\partial u}{\partial t} = \left( \frac{\partial u_1}{\partial t}, \frac{\partial u_2}{\partial t}, \frac{\partial u_3}{\partial t} \right) \text{ and } \frac{\partial u}{\partial s} = \left( \frac{\partial u_1}{\partial s}, \frac{\partial u_2}{\partial s}, \frac{\partial u_3}{\partial s} \right).$$

- Prove that the pull-back by  $u$  of the Riemannian area form on  $S$  coming from the metric induced by the inner product on  $\mathbb{R}^3$  is

$$\left\| \frac{\partial u}{\partial t} \times \frac{\partial u}{\partial s} \right\| dt \wedge ds.$$

- Let  $v$  be a vector field on  $\mathbb{R}^3$  and  $\omega = i_v(dx \wedge dy \wedge dz)$ . Prove that

$$u^*\omega = v \cdot \left( \frac{\partial u}{\partial t} \times \frac{\partial u}{\partial s} \right) dt \wedge ds,$$

and hence, in the sense of the standard vector calculus,

$$\int_u \omega = \iint_S v \cdot dS.$$