

MATH 209, MANIFOLDS II, WINTER 2006

Homework Assignment I: One-forms and integration,  
due Thursday 1/19-2006

1. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be given by  $y = x^3$ . Calculate  $f_*(\partial/\partial x)$  and  $f^* dy$ . Is  $f_*(\partial/\partial x)$  smooth?
2. Let  $f$  be a smooth function on a manifold  $M$  and  $(x_1, \dots, x_n)$  a system of local coordinates. Prove that in local coordinates

$$df = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i.$$

Remark. Here one has to be careful with definitions. In this problem we define the one-form  $df$  by  $df(v) = L_v f$ . The one-forms  $dx_1, \dots, dx_n$  are defined as the basis in  $T_x^*M$  dual to the basis  $\partial/\partial x_1, \dots, \partial/\partial x_n$  in  $T_x M$ . (Note that alternatively one could define  $dx_i$  by  $dx_i(v) = L_v x_i$ , where  $x_i$  is now thought of as a function defined on an open subset of  $M$ . It follows from the assertion of the problem that this definition is equivalent to the one through the dual basis.)

3. Let  $\alpha = x dy$  on the plane  $\mathbb{R}^2$  with coordinates  $(x, y)$ . Denote by  $S_R^1$  the circle of radius  $R > 0$  centered at the origin, oriented counter clockwise. Evaluate

$$\int_{S_R^1} \alpha.$$

Conclude from the result that  $\alpha$  is not exact, i.e., there exists no function  $f$  such that  $\alpha = df$ . More generally, let  $\gamma$  be a closed simple curve in  $\mathbb{R}^2$ . What is the geometrical meaning of  $\int_{\gamma} x dy$ ?

4. Let  $\alpha$  be a one-form on a smooth connected manifold  $M$ . Recall that  $\alpha$  is said to be exact if there exists a function  $f$  such that  $\alpha = df$ . Prove that

- $\alpha$  is exact  $\iff \int_{\gamma} \alpha = 0$  for any closed curve  $\gamma$ .

Remark. The part  $\implies$ ), which we did in class, is here only for the sake of completeness. You do not need to redo it. Here is a hint for part  $\impliedby$ ). The goal is to construct  $f$  such that  $\alpha = df$ . Fix a reference point  $x_0 \in M$ . For  $x \in M$ , set  $f(x) = \int_{\eta} \alpha$ , where  $\eta$  is curve connecting  $x_0$  to  $x$ . First show that the function  $f$  is well defined, i.e.,  $f(x)$  is really independent of the curve  $\eta$  connecting  $x_0$  and  $x$ , using the assumption that  $\int_{\gamma} \alpha = 0$ . Then prove that  $df = \alpha$ . (For instance, one can use local coordinates to this end.)

5. Problem 6-7 (Chapter 6) on page 152 of the textbook.